

# [Beautiful healthy legs]

# Aethoxysklerol®



Sclerotherapy is considered to be the method of first choice for the treatment of small calibre varicose veins.\*

Sklerosierungsmittel  
Sclerosing agent  
Agent sclérosant  
Sostanza sclerosante  
Medio esclerosante

\* Translated from  
German Association of the Scientific Medical Societies (AWMF) - Standing Guidelines Commission.  
"Sklerosierungsbehandlung der Varikose". Date 31.12.2018. Available at: <https://www.awmf.org/leitlinien/detail/ll/037-015.html>

Kreussler Pharma

POB 12 04 54 • D-65082 Wiesbaden • Rheingastr. 87-93 • D-65203 Wiesbaden  
Phone: +49 611 9271-0 • Fax: +49 611 9271-111  
[www.kreussler.com](http://www.kreussler.com) • [info@kreussler.com](mailto:info@kreussler.com)

**kreussler**  
PHARMA



# Aethoxysklerol®



maximum number of ampoules to reach the maximum dose of 2 mg polidocanol per kg bodyweight in a patient weighing

50 kg 60 kg 75 kg 90 kg

	2.5 mg/ml Polidocanol <sup>1</sup>	20	24	30	36				
	5 mg/ml Polidocanol <sup>1</sup>	10	12	15	18				
	10 mg/ml Polidocanol <sup>1</sup>	5	6	7.5	9	2			
	20 mg/ml Polidocanol <sup>1</sup>	2.5	3	3.5	4.5				
	30 mg/ml Polidocanol <sup>1</sup>	1.5	2	2.5	3				
Volume (ml) per injection					0.1 – 0.2 ml	0.1 – 0.3 ml	0.5 – 2.0 ml <sup>3</sup>	0.5 – 2.0 ml <sup>3</sup>	0.5 – 1.0 ml <sup>4</sup>

- Aethoxysklerol®** universell einsetzbar bei optimaler Wirkung und Verträglichkeit
- Aethoxysklerol®** universally applicable with an optimum of efficacy and tolerance
- Aetoxisclerol®** universellement applicable avec efficacité et tolérance optimales
- Atosisclerol®** utilizzo universale con un massimo di efficacia e tolleranza
- Etoxisclerol®** aplicación universal con efecto y tolerancia óptima

Availability of different concentrations depends on national marketing authorisation.  
Dosage according to the German SPC.

<sup>1</sup> Polidocanol, also known as Lauromacrogol 400, is produced by Kreussler and GMP-approved

<sup>2</sup> For spider veins, Aethoxysklerol 1% is administered only in central veins of spider veins

<sup>3</sup> In the first treatment session, only one injection of 0.5 – 1 ml should be given

<sup>4</sup> For sclerotherapy of haemorrhoidal disease, a total of 3 ml should not be exceeded during one treatment session; when treating an 11 o'clock haemorrhoid in men, the quantity injected must not exceed 0.5 ml